

**Induced Lorentz-Violating Chern-Simons Term in QED:
Uncovering Short Distance Interaction Terms in the Effective
Lagrangian without the Shadow of Regularization***

Lai-Him Chan

*Department of Physics and Astronomy, Louisiana State University, Baton Rouge,
Louisiana 70803-4001, USA
E-mail: chan@phys.lsu.edu*

We show that the correctly evaluated effective Lagrangian should include short-distance interaction terms which have been avoided under the protection of usual regularization and must be properly identified and reinstated if regularization is to be removed. They have special physical and mathematical significance as well as restoring gauge invariance and suppressing divergence in the effective Lagrangian. The rich structure of the short-distance interaction terms can open up challenging opportunities where the conventional regularization with rigid structure is unavailable and inappropriate. It becomes clear that gauge invariance is preserved with or without regularization and therefore there is no Lorentz-Violating Chern-Simons term in QED.

There has been a serious controversy generated by a series of recent papers whether a Lorentz-violating CPT-odd term in a charged fermion Lagrangian,

$$\mathcal{L} = \bar{\psi} \left[i\rlap{\not{\partial}} - m - Q\rlap{\not{A}}(x) - \gamma_5 \rlap{\not{b}} \right] \psi, \quad (1)$$

can induce a finite Lorentz-violating term $\frac{1}{2}\epsilon_{\mu\alpha\beta\nu}k^\mu F^{\alpha\beta}A^\nu$ to the Lagrangian in the electromagnetic sector. The values of k_μ from theoretical calculations varies from 0, $\frac{3}{8\pi^2}Q^2b_\mu$, $\frac{1}{8\pi^2}Q^2b_\mu$ to completely undetermined^{1,2,3,5,6,7}.

The effective action resulted from integrating out the fermion field in Eq.(1) is given by,

$$\Gamma_{eff} = -i Tr \ln \left[i\rlap{\not{\partial}} - Q\rlap{\not{A}}(x) - m - \gamma_5 \rlap{\not{b}} \right], \quad (2)$$

The Tr here denotes both the trace tr_γ in the Dirac spinor space and the trace tr_x in the coordinate space. Using the following method to evaluate tr_x ⁸,

$$\begin{aligned} tr_x f(i\partial - QA(x)) &= \int d^4x \lim_{x' \rightarrow x} \langle x' | f(i\partial - QA(x)) | x \rangle = \int d^4x \lim_{x' \rightarrow x} f(i\partial - QA(x)) \delta(x - x') \\ &= \int d^4x \int \frac{d^4p}{(2\pi)^4} \lim_{x' \rightarrow x} f(i\partial - QA(x)) e^{-ip \cdot (x - x')} = \int d^4x \int \frac{d^4p}{(2\pi)^4} f(i\partial + p - QA(x)), \end{aligned} \quad (3)$$

*Talk presented at the Second Meeting on CPT and Lorentz Symmetry CPT'01, August 15-18, 2001, Bloomington, Indiana.

we obtain the effective Lagrangian in a compact form from Eq.(2)

$$\mathcal{L}_{eff} = -i \int \frac{d^4 p}{(2\pi)^4} \text{tr}_\gamma \ln G^{-1}(p + i\partial, x), \quad (4)$$

where $G(p + i\partial, x) = [\not{p} - m + i\not{\partial} - Q\not{A}(x) - \gamma_5 \not{b}]^{-1}$ and p is the fermion loop momentum.

When gauge invariant regularization is used, the effective Lagrangian is gauge invariant and naturally $k_\mu = 0$ ^{1,4}. It has been argued that if the gauge noninvariant anomalous contributions is finite without regularization, then the role of a gauge-invariant regularization is no longer regulating the divergent quantities but rather imposing gauge invariant^{2,5,6}. The option of removing gauge-invariant regularization should be kept open, in this case as in the chiral anomaly calculation. However, without the protection of regularization, it is not clear to what degree short-distance dynamics has been or can ever be properly defined. Therefore k_μ may not be determined⁷.

To uncover possible short-distance terms, we have to make precise calculation of the effective Lagrangian Eq.(4)⁶. Without the protection of regularization, the intrinsic quantum incompatibility, the noncommutativity between the photon fields and the space-time derivative, becomes relevant⁶. The power expansion of the photon fields and the power expansion of the derivative (small momentum) expansion become incompatible when the coefficients of lower power of A become singular. This difficulty can be only circumvented by expanding the effective Lagrangian in powers of $\not{A} = i\not{\partial} - Q\not{A}(x)$ and $\gamma_5 \not{b}$. We obtain the Chern-Simons Lagrangian⁶,

$$\mathcal{L}_{eff}^{CS} = 4i \epsilon_{\mu\alpha\beta\nu} b^\mu \Pi^\alpha \Pi^\beta \Pi^\nu \int \frac{d^4 p}{(2\pi)^4} \frac{m^2}{(p^2 - m^2)^3} = \frac{i}{16\pi^2} Q^2 F^{\alpha\beta} A^\kappa \epsilon_{\lambda\alpha\beta\nu}.$$

In order to compare our result with the calculation basing on the Feynman diagram approach, we compute the vacuum polarization tensor,

$$\begin{aligned} \Pi^{\mu\nu}(q) = & \int \frac{d^4 p}{(2\pi)^4} \text{tr}_\gamma \left\{ \gamma^\mu \frac{i}{\not{p} - m - \gamma_5 \not{b}} \gamma^\nu \frac{i}{\not{p} + \not{q} - m - \gamma_5 \not{b}} \right. \\ & \left. + i q_\alpha \frac{\partial}{\partial p^\alpha} \int_m^\infty dM \left[\gamma^\mu \left(\frac{i}{\not{p} - M - \gamma_5 \not{b}} \right)^2 \gamma^\nu \frac{i}{\not{p} + \not{q} - M - \gamma_5 \not{b}} \right] \right\} \end{aligned} \quad (5)$$

which is written in terms of the standard one-loop vacuum polarization Feynman amplitude with b_μ -exact fermion propagators² plus a b_μ -exact correction from the noncommutativity of the operators A and ∂ ⁶. This correc-

tion gives precisely the additional contribution $-\frac{1}{32\pi^2}Q^2b_\mu$ to k_μ to be combined with the result from previous calculation² of the first term to yield $k_\mu = \frac{3}{32\pi^2}Q^2b_\mu - \frac{1}{32\pi^2}Q^2b_\mu = \frac{1}{8\pi^2}Q^2b_\mu$.

By removing regularization, gauge dependent short-distance interaction terms, in the form of momentum integral of total momentum derivative not expressible in term of conventional Feynman diagrams, are exposed and can make finite contributions to the effective Lagrangian. The finite nongauge-invariant anomalous contributions are due to the short-distance singularity and sensitively depend on the path how x' approaches to x . In Eq.(3), tr_x should be performed symmetrically with respect to x and x' , However, the reduction from Eq.(2) to Eq.(4) by Eq.(3) is explicitly asymmetric in x and x' because the operator $i\partial - QA(x)$ is not diagonal in x and cannot be diagonalized. It appears that there is no simple way to escape from regularization. However, there is a very good reason and a solution for this problem.

In the semiclassical approach, the particle field is divided into a classical part which satisfies the classical field equation and the quantum part which obeys the quantum canonical commutation or anticommutation relations. In the path integral approach, the effective action is obtained by functionally integrating out the quantum fluctuation due to the quantum part. Likewise, in the evaluation of tr_x , the pair of space-time canonical operators in Eq.(4), $\{p + i\partial, x\}$, is the sum of the quantum part and the classical part. The pair $\{i\partial, x\}$ obeying the same canonical commutation relation is obviously the quantum part and $\{p, 0\}$ diagonal in p must be the classical part. Such decomposition is completely opposite to the original semiclassical approach that one would expect. Namely, x , being the space-time coordinates of the background field, should be classic and the loop-momentum p representing the integration of quantum fluctuation should be the quantum part. Therefore the decomposition of the canonical pair in Eq.(4) should have been $\{p, x - i\frac{\partial}{\partial p}\}$. The two decompositions are related by a gauge covariant unitary transformation $e^{\Pi \cdot \frac{\partial}{\partial p}}$ in the direct product space of x and p ⁹. The expansion to this larger space allows the diagonalization of x .

Therefore the correct effective Lagrangian with interaction $\chi(x)$ in any dimension N should be,

$$\mathcal{L}_{eff} = -i \int \frac{d^N p}{(2\pi)^N} tr_\gamma \left[e^{-\Pi \cdot \frac{\partial}{\partial p}} \right] \ln (\not{p} + \not{\Pi} - m - \chi(x)) \left[e^{\Pi \cdot \frac{\partial}{\partial p}} \right], \quad (6)$$

which is gauge invariant and diagonal in x but not in p . The corresponding propagator is,

$$G(p, \Pi) = \left[e^{-\Pi \cdot \frac{\partial}{\partial p}} \right] \frac{i}{\not{p} + \not{\Pi} - m - \chi(x)} \left[e^{\Pi \cdot \frac{\partial}{\partial p}} \right]. \quad (7)$$

In Eq.(6), the expansion of $e^{\Pi \cdot \frac{\partial}{\partial p}}$ gives no contribution. The expansion of $e^{-\Pi \cdot \frac{\partial}{\partial p}}$ yields correction terms proportional to the total derivative of p . They have been safely discarded in conventional calculation with regularization. They must be reinstated if regularization is not used. Those correction terms can be interpreted as short-distance gauge interaction and they are just as well defined as conventional minimal gauge interaction in the Lagrangian by the requirement for maintaining gauge invariance and minimizing of the degree of divergence. Without regularization, it takes the coordination of both the long distance interaction and short distance interactions to fulfill that requirement. There is also a simple physical interpretation. Photon emission is accompanied not only with the change of momentum of charge classical momentum but also with the change of loop momentum due the quantum fluctuation at short distance. The Green's function in Eq.(7) is similar in form to the Schwinger's point splitting method of bilinear current. Here, the short distance singularity due to two field operators at the same space-time point is averaged out due to quantum fluctuation rather than classical average. $e^{(i\partial - eA) \cdot \frac{\partial}{\partial p}}$ is a covariant displacement operator to displace x by $i \frac{\partial}{\partial p}$.

$$e^{(i\partial - eA) \cdot \frac{\partial}{\partial p}} \psi(x) = \psi(x + i \frac{\partial}{\partial p}) \quad (8)$$

Such operator can easily be constructed for any gauge symmetry and any particle fields.

We shall use the schwinger model, the 1+1 dimensional massless QED, to test the validity of the short-distance interaction terms in the effective Lagrangian. We rewrite Eq.(6) with $N=2$, $m=0$, and $\chi(x)=0$ in a form ready to expand in power series of A .

$$\mathcal{L}_{eff} = - \int \frac{d^2 p}{(2\pi)^2} \left[e^{-(i\partial - eA) \cdot \frac{\partial}{\partial p}} \right] \int_0^\infty dm \operatorname{tr}_\gamma \frac{1}{\not{p} + i\not{\partial} - e\not{A} - m}. \quad (9)$$

We have dropped the factor $e^{\Pi \cdot \frac{\partial}{\partial p}}$ because it has no contribution. The vacuum polarization can be calculated directly without any regularization,

$$\Pi^{\mu\nu}(p) = \frac{e^2}{\pi} \left(\frac{1}{2} g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) - \frac{e^2}{2\pi} g_{\mu\nu} + \frac{e^2}{\pi} g_{\mu\nu} = \frac{e^2}{\pi} \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right). \quad (10)$$

The first term corresponds to the contribution from conventional calculation without the exponential correction factor⁵. The second term comes purely from the short distance correction factor. The last term is from the mixed contribution. The net result recovers the correct solution with photon mass

$M^2 = \frac{e^2}{\pi}$. Therefore the short distant interaction terms from the gauge covariant unitary transformation has indeed fulfilled its promised role of restoring gauge invariance as well as contributing the essential correction to the unregulated vacuum polarization.

The CPT and Lorentz-violating effective Lagrangian corresponds to the Eq.(6) with $N = 4$ and $\chi(x) = \gamma_5 \not{x}$. Since the short-distance interaction terms has restored gauge invariance, It is obvious that there is no induced Lorentz-Violating Chern-Simons term.

We can now safely abolish the regularization barrier and reclaim the territory of short distance interaction in quantum field theory. This new degree of freedom as viewed by individual Feynman amplitudes appears to be chaotic. However in the background field hybrid effective Lagrangian approach, we have the benefit of viewing the generating functional of the complete set of Feynman amplitudes and a new prospective begin to emerge. We have demonstrated that not only short-distance interaction can be derived to replace the effect of usual regularization but the process can have real mathematical and physical significance. The short-distance interaction can be considered as the quantum images of the long-distance interaction in the classical Lagrangian. They must be chosen in tandem to maintain gauge symmetries and to minimize the degree of divergence in higher order perturbation calculations. It will be a great challenge to apply this new tool to the cases where the conventional regularization is unavailable or inappropriate.

References

1. S. Coleman and S. L. Glashow, Phys. Rev. **D59** 116008 (1999).
2. D. Colladay and V.A. Kostelecky, Phys. Rev. **D 58**, 116002 (1998). R. Jackiw and V. A. Kostelecký, Phys. Rev. Lett. **82**, 3572 (1999).
3. M. Pèrez-Victoria, Phys. Rev. Lett. **83**, 2518 (1999); J. M. Chung, Phys. Lett. **B461**, 138 (1999); M. Chaichian, W. F. Chen, and R. González Felipe, Phys. Lett. **B503**, 215 (2001)
4. G. bonneau, Nucl. Phys. **B593**, 398 (2001); C. Adam and F. R. klinkhamer, Phys. Lett. **B513**, 245 (2001)
5. R. Jackiw, Int. J. Mod. Phys. **B14**, 2011 (2000).
6. Lai-Him Chan, hep-ph/9907349
7. W. F. Chen, Phys. Rev. **D60** 085007 (1999); J. M. Chung, Phys. Rev. **D60** 127901 (1999).
8. Lai-Him Chan, Phys. Rev. Lett. **54**, 1222 (1985); **56**, 404E (1985); **55**, 21 (1985); **57**, 1199 (1986); Phys. Rev. **D55**, 6223 (1997).
9. M. K. Gaillard, Nucl. Phys. **B268**, 669 (1986).